

Perianal Skin Tag

Operations in or around anus will be made, if big hemorrhoides, loose skin after hemorrhoides, abscesses, or treatment for chronic crack in the anus area, will not heal on medical treatment.

Before the operation:

3 – 4 days before the operation, start taking 2 – 3 tablets Magnesia 500 mg in the evening, to make sure stools become soft. Remember to drink plenty of liquids. Put on the Emla/Tapin creme around the anus 1-2 hours before the operation. You will be able to go home after the operation, but no biking.

Operation:

The operation will be done in local anesthesia, which is installed via a plug, in the hemorrhoid, or the around area in the anus.

After the operation:

It's a very good idea to lay down on the very first day after the operation, course it will facilitates the pains. Also the very first 2 – 3 days after, can be pain full, why the Magnesia tablets are important. You can advantageously have some Paracetamol tablets, or Ibuprofen, unless other agreements.

Dressing can be removed the day after the operation. The surgical wounds are best to clean by rinse off, with a hand shower twice a day, and after toilet visits, until the operation area will be painless. Optionally you can put on some cold covers, to stop the bleedings, or to reduce the pain. Stitches should not be removed.

We will check it up after 10 – 12 days after the operation.

Sickleave: can be necessary for about 3 – 5 days.

Complications: There may be some bleeding in the following days, major bleeding are very rarely, as well infections.

Seek medical advice:

In case of major bleeding or infection, please contact the surgical doctor, doctor on call, or the emergency room right away.